

Composition: Systematic List of Species

<i>Betula ashleyi</i>		26
<i>Pinus ponderosoides</i>		23
<i>Mahonia reticulata</i>		23
<i>Pinus balfouroides</i>		21
winged seeds	16	
fascicles	4	
branchlet		
<i>Populus pliotremuloides</i>		18
<i>Populus payettensis</i>		17
<i>Larix churchillensis</i>		14
winged seeds		
<i>Mahonia macginitiei</i>		14
<i>Prunus treasheri</i>		12
<i>Amelanchier desatoyana</i>		12
<i>Carya bendirei</i>		11
<i>Populus eotremuloides</i>		10
<i>Lyonothamnus parvifolius</i>		10
<i>Quercus wislizenoides</i>		9
<i>Rosa harneyana</i>		9
<i>Nymphaeites nevadensis</i>		8
leaves	3	
root scars	5	
<i>Chrysolepis sonomensis</i>		8
<i>Acer oregonianum</i>		7
samaras		
<i>Heteromeles desatoyana</i>		7
<i>Prunus chaneyii</i>		6
<i>Salix ovyhecana</i>		6
<i>Salix laevigatoides</i>		5
<i>Fraxinus eastgatensis</i>		5
leaflet		
samaras	4	
<i>Cedrela trainii</i>		4
<i>Salix desatoyana</i>		3
<i>Ribes webbi</i>		3
<i>Crataegus middlegatei</i>		3
<i>Eugenia nevadensis</i>		3
<i>Hydrangea bendirei</i>		3
<i>Ribes barrowsii</i>		3
<i>Abies concoloroides</i>		2
winged seeds		
<i>Abies laticarpus</i>		2
winged seeds		
<i>Ribes bonhamii</i>		2
<i>Amorpha stenophylla</i>		2
<i>Symphoricarpos wassukana</i>		2
<i>Alnus latahensis</i>		2
<i>Chamaecyparis cordilleriae</i>		1
<i>Acer medianum</i>		
<i>Betula idahoensis</i>		
<i>Ribes stanfordianum</i>		
<i>Cercocarpus ovatifolius</i>		
<i>Chamaebatia nevadensis</i>		
<i>Prunus moragensis</i>		
<i>Robinia bisonensis</i>		
<i>Acer trainii</i> (samara)		
<i>Acer tyrrellii</i> (samara)		1
<i>Vaccinium sopheroides</i>		1

Total: 8,276

*These fossils occur along strike for about 40 m and are distributed stratigraphically for about 1 m